



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

wave of November 1 and 2, 1911, brought the ducks south almost by the millions. The Eider was shot by a Doctor Page of Sioux Falls and is now in the possession and displayed in the place of business of William Sweet of this city.—S. S. VISHNER, *University of South Dakota, Vermillion, S. D.*

Records from Nippinickett Pond, Bridgewater, Mass.—The captures of the following ducks on Nippinickett Pond seem worthy of record. European Widgeon, *Mareca penelope*, Nov. 7, 1902. Shot by A. C. Dyke.

Two European Widgeons, *Mareca penelope*, Oct. 22, 1910, from a flock of 4 birds. Shot by Harry P. Sturtevant.

European Widgeon, *Mareca penelope*, Oct. 23, 1910. Shot by A. C. Dyke.

Hutchin's Goose, *Branta canadensis hutchinsi*, Oct. 8, 1910, from a flock of six birds, the other five being Canada Geese. Shot by Messrs. Frank C. Drake and Irving A. Hall.

All the specimens were obligingly identified by Mr. Wm. Brewster of Cambridge.—ARTHUR C. DYKE, *Bridgewater, Mass.*

Sora Rail (*Porzana carolina*) in New Mexico.—On the morning of August 27, 1912, at State College, New Mexico, I picked up, dead, an immature Sora Rail that had flown against a telephone wire and broken its neck. On the night of August 23, 1912, about 10 o'clock P. M., a flock of birds flew over me on the mesa, headed northeast—away from the Rio Grande. The size of the birds and their manner of flight, as seen by the moonlight, in conjunction with the finding of the above specimen, lead me to believe these birds were of the same species.

To my knowledge, this is the first specific record of this species in the state, certainly in this locality. There has been an unusual amount of rainfall in this region during the month of August, which may account for their presence here at this time, for our mesa country offers no inducements to them naturally.—D. E. MERRILL, *State College, N. M.*

Early Occurrence of the White-rumped Sandpiper (*Pisobia fuscicollis*) in Maine.—A male White-rumped Sandpiper in nuptial plumage was secured on April 27, 1912, at Scarborough Maine. The bird was alone.

The only other instances of its occurrence in the state in spring, known to me, are those recorded by Mr. Nathan Clifford Brown, May 30, 1881, and May 29, 1882.¹ Both of these recorded by Mr. Brown are in the collection of the Portland Society of Natural History, and the one here recorded in that of the writer.

The bird according to Mr. W. W. Cooke is rare on the Atlantic coast of the United States, in spring, north of Virginia,² and its occurrence at this

¹ Proc. Portland Society of N. H., II, p. 27.

² Bull. 35, U. S. Biol. Survey, p. 38.